



ISSN: 2250-2823

HortFlora

Research Spectrum

Volume 6, Issue 2
June 2017

An International
JOURNAL
Peer Reviewed
NAAS Rating : 3.78



BIOSCIENCES & AGRICULTURE ADVANCEMENT SOCIETY

www.hortflorajournal.com

ISSN: 2250-2823



HortFlora

Research Spectrum

Peer Reviewed

An International

JOURNAL

International Impact

Index Copernicus Value (ICV) : 27.39; Global Impact Factor (GIF) : 0.471
InfoBase Index (IBI) Factor : 2.8; New Journal Impact Factor (NJIF) : 2.14
Open Academic Journals Index (OAJI) Impact Factor : 0.201

Volume 6, Issue 2

June 2017

Indexed / Abstracted in :

- Index Copernicus International, Poland
- Indian Science Abstracts
- CAB Abstracts
- CABI Full text
- CiteFactor
- OAJI.net
- I2OR
- Spice Bibliography
- InfoBase Index
- Google Scholar
- Research Bib
- ICRISAT InfoSAT
- getCited
- JournalIndex.net
- ISRAJIF
- NJIF

NAAS Rating : 3.78

UGC Approved
(Jour. No. 45792)



BIOSCIENCES & AGRICULTURE ADVANCEMENT SOCIETY

www.hortflorajournal.com

Date of Publication : 02-07-2017



CONTENTS

1. Role of Major Nutrients (NPK) in Flower Crops : A Review	G. Madhuri, B. Babu Rao and P. Neelima	69-76
2. Biology and Eco-friendly Management of 28-Spotted Lady Beetle (<i>Henosepilachna vigintioctopunctata</i> Fabr.)—A Serious Pest of Brinjal	Arti Katiyar, Vijai Kumar and K.D. Verma	77-81
3. An Economic Assessment of Banana Production in Bhagalpur district of Bihar	Priyanka Kumari, Nasim Ahmad and M.K. Wadhvani	82-87
4. Floral Biology Studies in Guava (<i>Psidium guajava</i> L.) Genotypes	Banoth Shiva	88-92
5. Optimization of Protocol for Minimizing <i>in-vitro</i> Contamination of Embryo Cultured Peach (<i>Prunus persica</i> L.)	A.S. Sundouri, Harminder Singh, Anirudh Thakur and Dimpy Raina	93-96
6. Physico-Chemical and Morphological Evaluation of Guava (<i>Psidium guajava</i> L.) Genotypes under <i>Tarai</i> Conditions.	Jitendra Kumar, Rajesh Kumar, Shailesh Tripathi and Vijay Pratap Singh	97-101
7. Prevalence of Citrus Diseases in Different Agro-Climatic Zones of Punjab	Anita Arora	102-105
8. Physico-Chemical Changes During Growth, Maturity and Ripening in Ber (<i>Ziziphus mauritiana</i> Lamk) cv. Gola and Chhuhara	V.K. Asati, K. P Singh and A. K. Goswami	106-109
9. Growth and Yield of Aloe (<i>Aloe barbadensis</i> M.) as Influenced by Harvest Dates	Bijit Kumar Saud and Urbashi Hazarika	110-113
10. Effect of Different Organic Manures and Bio-fertilizers on Growth and Yield of Okra and Nutrient Uptake	V. V. Appa Rao, Sanjay Singh and S. Raja	114-118
11. Performance of Tuberose (<i>Polianthes tuberosa</i> L.) Cultivars Grown for Spikes under Western Uttar Pradesh Conditions	Virendra Pal, K.P. Singh, Ashish Tyagi, Anant Kumar and Omvir Singh	119-121
12. Fruit Production and Biochemical Aspects of Seeds of <i>Euryale ferox</i> Salisb. under <i>ex-situ</i> Conditions	Amit Kumar, I.S. Singh, Ramesh Kumar, V. N. Jha, A. K. Thakur and Anil Kumar	122-124
13. Effect of Stage of Maturity and Method of Drying on Bio-Chemical and Organoleptic Attributes of Ber (<i>Ziziphus mauritiana</i> Lamk.) Fruits	V.K. Asati, K. P Singh and A. K. Goswami	125-127
14. Response of Bulb Size on Vegetative Growth, Flowering and Bulb Production in Tuberose (<i>Polianthes Tuberosa</i> L.) cv. Phule Rajani	Krishan P. Singh, Tarak N. Saha and P. Hollajer	128-130
15. Effect of NPK on Plant Growth and Flowering of <i>Amaryllis belladonna</i>	Rohit Kumar Sharma, S. Saravanan and Sunita Kumari	131-133
16. Effect of Mulching on the Growth and Yield of Onion (<i>Allium cepa</i> L.)	S. K. Singh, Sanjay Kumar ² and Subash Chand	134-136

ABSTRACTS

NAAS Rating : 3.78

www.hortflorajournal.co

HortFlora Research Spectrum, 6(2) : (June 2017)

ISSN : 2250-2823



1. Role of Major Nutrients (NPK) in Flower Crops : A Review

G. Madhuri¹, B. Babu Rao² and P. Neelima³

¹Deptt. of Floriculture and Landscaping, OUAT, Orissa, India

²Deptt. of Vegetable Science, Dr. Y. S. R. H. U, V. R. Gudem, Andhra Pradesh

³Deptt. of HC & RI, Dr. Y. S. R. H. U, V. R. Gudem, Andhra Pradesh

*Corresponding Author's E-mail: neelimapalagani@yahoo.com

ABSTRACT : Vibrant horticulture is dependent upon many factors, among which proper and judicious soil nutrient management practices comes first. There are seventeen elements are known to be essential for plants. Among which nitrogen, phosphorous and potassium plays vital role and they are considered as major plant nutrients. Among horticultural crops, floriculture is growing at an alarming rate both in open and protected cultivation. Proper growth and development of the plant is ultimately effected by regular nutritional management. Importance of major plant nutrients and their dose for better growth and yield for various flower crops viz. gladiolus, rose, gerbera, golden rod, cymbidium, chrysanthemum, calendula, China aster, anthurium, tulip, marigold, tuberose, gaillardia, larkspur, spider lily etc. has been reviewed for better understanding and will be helpful for present crop cultivation and future research purpose.

Published in : HortFlora Research Spectrum, 6 (2) : 69-76 (June 2017)

1. Biology and Eco-friendly Management of 28-Spotted Lady Beetle (*Henosepilachna vigintioctopunctata* Fabr.)—A Serious Pest of Brinjal

Arti Katiyar^{1*}, Vijai Kumar² and K.D. Verma³

¹ICAR- Indian Institute of Pulses Research, Kanpur (U.P.)- 208 024, India

²Deptt. of Horticulture, CSSS (PG) College, Machhra, Meerut

³Deptt. of Entomology, A.S. College, Lakhaoti, Bulandsahar

*Corresponding Author's E-mail: artikatiyar25@gmail.com

ABSTRACT : Brinjal (*Solanum melongena* L.) is an important Solanaceous crop of subtropics and tropics. Several biotic and abiotic factors contribute in lowering the yield in brinjal. Among various biotic factors, insect pests are important which greatly affect the quality and productivity of brinjal crop through inflicting a direct damage. 28-Spotted lady beetle, *Henosepilachna vigintioctopunctata* (F.) (Coleoptera: Coccinellidae) is one of the major pest of egg plant. The larvae and adults scrape the green matter from brinjal leaves and cause damage up to 80%. The morphology and biology of the pest is described under this study. Among the various components of management, *Aspergillus flavus* and *Bacillus thuringiensis* are effective against different stages of *H. vigintioctopunctata*. The predator reduviid bug, *Rhinocoris fuscipes* (Fab.) is quite efficient. The various plant products are found to be effective against this pest. The repellent and antifeedant properties of Neem, Mahua and Groundnut cakes have also been established.

Published in : HortFlora Research Spectrum, 6 (2) : 77-81 (June 2017)

3. An Economic Assessment of Banana Production in Bhagalpur district of Bihar

Priyanka Kumari¹, Nasim Ahmad^{1*} and M.K. Wadhwani²

¹Department of Agricultural Economics, Dr. Rajendra Prasad Central Agricultural University, Pusa Samastipur-848 125 (Bihar),

²Department of Agricultural Economics, Bihar Agricultural University, Sabour, Bhagalpur

*Corresponding author's E-mail: nasim.rau@gmail.com

ABSTRACT : The study was conducted on primary data, collected from a sample of 60 banana growers, consisted of 26 (43 %), 25 (42 %) and 9 (15 %) marginal & small, semi-medium and medium & large category. The data were collected from the sample growers with the help of specifically prepared pre-tested schedules through Survey Method by interviewing them, selected through Multi-Stage Sampling Technique from a cluster of 3 villages of Nawgachia block of Bhagalpur district. The study revealed that per ha average total cost of cultivation of banana var. robusta was found ranging between ₹ 1, 00,566.45 on marginal & small farms to ₹ 95, 294.42 on medium & large farms. The variable costs included material cost (72.55 %) and labour cost (23.60 %). The material cost was found highest (₹ 72364.71 per ha) on marginal & small farms. The average labour cost was estimated as ₹ 22512.93 per ha. The average cost of irrigation and fertilizer constituted 35.59 per cent and 16.01 per cent of total cost, respectively while labour cost constituted of 23.60 per cent. The yield (number of bunches per ha) was found ranging from 2768 to 2847 among three categories of farms and the average price received by the banana growers was ₹ 129.91 per bunch. The average gross income has been estimated as ₹ 3, 64,007.87 per ha, which was ranging in ascending order to size of farms; i.e., higher on large farms and less on marginal farms. The “return to total cost ratio” was estimated at 3.78, which was also in ascending order to size group.

Published in : HortFlora Research Spectrum, 6 (2) : 82-87 (June 2017)

4. Floral Biology Studies in Guava (*Psidium guajava* L.) Genotypes

Banoth Shiva*

Div. of Fruits and Horticultural Technology,

ICAR-Indian Agricultural Research Institute, New Delhi-110 012, India

*Corresponding Author's E-mail: banothshivaari@gmail.com

ABSTRACT : This experiment was carried out to obtain information on flowering pattern of *Psidium*, so that suitable species or varieties can be used in future breeding programme for their hybridization work to plan and develop crosses. The results revealed that cultivar Allahabad Safeda was earliest to bloom (24th April) followed by Arka Amulya, Black guava, Hissar Surkha, Lalit and Snow White (25th April), whereas L-49 (Sardar) was last to flower (10th May). Flowering duration among guava genotypes differed significantly and it ranged between 39 to 52 days. The maximum flowering duration was observed in Lalit and Shweta (52 days) followed by Hissar Surkha (51 days), Sasni collection (51 days) and Arka Amulya (50 days), while, the shortest flowering duration was recorded in Hafsi Red (39 days). Thus, from the above findings it was noticed that the variation between the genotypes may be attributed to the differences in the genetic makeup of these genotypes. Observation and knowledge of such traits can guide a programme of genetic improvement, especially those pursuing derivation of varieties through hybridization programme. Also trees can possibly be exploited by cultural manipulation to flower and fruit when desired.

Published in : HortFlora Research Spectrum, 6 (2) : 88-92 (June 2017)

5. Optimization of Protocol for Minimizing *in-vitro* Contamination of Embryo Cultured Peach (*Prunus persica* L.)

A.S. Sundouri¹, Harminder Singh², Anirudh Thakur² and Dimpy Raina³

¹Division of Fruit Science, Sher-e-Kashmir University of Agricultural Sciences & Technology of Kashmir, Shalimar, 190 025

²Department of Fruit Science, Punjab Agricultural University, Ludhiana, Pb., 141 004.

³KVK, Ferozepur, Punjab Agricultural University, Ludhiana, Pb., 141 004.

*Corresponding Author's Email: asundouri@gmail.com

ABSTRACT : Micro propagation is a rapid propagation technique for raising any kind of plant but it encounter with the utmost problem of contaminations by different sources such as bacteria, fungi and many more which reduces their productivity and completely prevents their successful cultivation. The present study was an attempt to minimize the contamination occurred in the embryo cultured peach by using different surface sterilants to obtain successful plantlet. Among the different sterilants cetrimide 5% for 15 minutes followed by carbandazium 2% for 5 minutes was the most effective treatment which showed maximum effect against contaminations. However, the sterilant 0.1% HgCl₂ for 10 minutes and ethanol for 70% for 3 minutes duration gave maximum uncontaminated plantlet with low survival rate.

Published in : HortFlora Research Spectrum, 6 (2) : 93-96 (June 2017)

6. Physico-chemical and Morphological Evaluation of Guava (*Psidium guajava* L.) Genotypes under Tarai Conditions

Jitendra Kumar*, Rajesh Kumar, Shailesh Tripathi and Vijay Pratap Singh

Department of Horticulture, College of Agriculture, Govind Ballabh Pant University of Agriculture & Technology Pantnagar (India)-263 145

*Corresponding Author's E-mail: jkumar_hort@outlook.com

ABSTRACT : A field experiment was conducted on different guava genotypes to evaluate the physico-chemical and morphological properties of guava. Different guava genotypes performed significantly different. MPUAT Sel-1 exhibited maximum fruit diameter, fruit size, fruit weight, fruit volume and total sugars percentage. While, Kayamganzi exhibited maximum fruit length and reducing sugars. Allahabad Safeda showed maximum length : diameter ratio. RCGH - 11 showed maximum specific gravity. RCGH-1 showed maximum TSS, titratable acidity, non-reducing sugars, sugar : acid ratio, plant height, plant canopy spread and stem girth. Sangareddy showed maximum ascorbic acid content. Other traits showed inferior performance among the evaluated guava genotypes.

Published in : HortFlora Research Spectrum, 6 (2) : 97-101 (June 2017)

7. Prevalence of Citrus diseases in Different Agro-climatic Zones of Punjab

Anita Arora*

Department of Fruit Science, Punjab Agricultural University, Ludhiana-141004, Punjab, India

*Corresponding Author's E-mail: anitapau@pau.edu

ABSTRACT : Surveys revealed that citrus foot rot/gummosis, fruit drop/stem-end rot and die-back were the most widespread diseases prevalent in Punjab. Other commonly occurring diseases were bacterial canker, citrus greening, citrus ring spot and sooty mould. Citrus foot rot/gummosis occurred in trees of all age groups across the agro-climatic zones. Population of *Phytophthora nicotianae* showed an increasing trend from July onwards and reached maximum in the month of September. Pathological fruit drop during the months of September and October was the most detrimental. Percent intensity of bacterial canker was very low in old orchards; whereas young orchards and nurseries in all the three agro-climatic zones were having low to moderate disease intensity. Citrus greening was more prevalent in the central and sub-mountainous regions, particularly on older trees. Moderate incidence of citrus ring spot was observed in arid-irrigated and central zones.

Published in : HortFlora Research Spectrum, 6 (2) : 102-105 (June 2017)

8. Physico-chemical changes during Growth, Maturity and Ripening in Ber (*Ziziphus mauritiana* Lamk) cv. Gola and Chhuhara

V.K. Asati¹*, K. P Singh¹ and A. K. Goswami²

¹Department of Horticulture, A. S. (P.G.) College, Lakhaoti, Distt. Bulandshahr, C.C.S. University Meerut (U.P.)

²Division of Fruits & Horticultural Technology, IARI, New Delhi.

*Corresponding Author's Email : drvinodasati@gmail.com

ABSTRACT : A study was conducted to evaluate the physical and chemical changes occurring during the various stages of fruit growth and development of 12-year-old Ber (cv. Gola and Chhuhara) trees. The ber fruit had a growth pattern of double sigmoid type. Fruit size rapidly increased from fruit set to 30th Nov. and comparatively slow from 30th Nov. to 30th Dec., thereafter it was increased upto 20th March. The fruit size remained unchanged after 20th March. Shape index was higher in initial stage and decreased continuously during the growth. The changes in fruit weight were almost similar to that of fruit length or diameter. The fruit weight increased with the advancement of maturity. The specific gravity of fruit showed a decreasing trend during the early stages and increased continuously until full fruit maturity. Pulp stone ratio was a slow and gradually increased after fruit set, followed by a sharp increase. The total soluble solids was low in early stage and it increased very much during later stages of fruit development. The per cent acidity was gradually increased up to 15th Feb. in cultivar Gola and upto 1st March in Chhuhara respectively, and it declined thereafter. Ascorbic acid content gradually increased from fruit set to maturity.

Published in : HortFlora Research Spectrum, 6 (2) : 106-109 (June 2017)

9. Growth and Yield of Aloe (*Aloe barbadensis* M.) as influenced by Harvest Dates

Bijit Kumar Saud* and Urbashi Hazarika

Department of Horticulture, College of Agriculture, Assam Agricultural University, Jorhat-785 013, India

*Corresponding Author's E-mail: bijit1969@rediff.com

ABSTRACT : To investigate the growth and yield of Aloe (*Aloe barbadensis* M.) in open condition as influenced by harvest dates, an experiment was carried out at the Medicinal and Aromatic Plants (MAP) Block, Experimental Farm, Department of Horticulture, Assam Agricultural University in Jorhat during 2015-2016. A total of four treatments with five replications were laid out in a Randomized Block Design. The treatments were comprised of: T₀- Harvesting at an interval of two months (recommended practice), T₁- Harvesting at one and a half months interval (after first harvest), T₂- Harvesting at two and a half months interval (after first harvest) and T₃- Harvesting at three and a half months interval (after first harvest). The results showed that the different harvest intervals had significant effect on the growth and yield of Aloe. The maximum values for all the growth and yield parameters were observed in T₃ [Harvesting at three and a half months interval (after first harvest)] treatment whereas the minimum values were observed in T₁ [Harvesting at one and a half months interval (after first harvest)] treatment. Overall, Aloe leaves harvested at three and a half months interval resulted in higher productivity and maximum benefit in subtropical climate of Jorhat, Assam.

Published in : HortFlora Research Spectrum, 6 (2) : 110-113 (June 2017)

10. Effect of different organic manures and bio-fertilizers on growth and yield of okra and nutrient uptake

V. V. Appa Rao*, Sanjay Singh and S. Raja

Central Horticultural Experiment Station, (ICAR-CIAH), Vejalpur- 389 340, Panchamahals, Gujarat.

*Corresponding Author's E-mail: vvapparaoyasu@gmail.com

ABSTRACT : An experiment was conducted in okra [*Abelmoschus esculentus* (L.) Moench.] with different organic manures like farm yard manure, vermi compost, poultry manure and sheep manure with biofertilizer combinations like *Azotobacter chroococcum* and *Azospirillum brasilense* with four replications in randomized block design during kharif season on an ustachrept soil of Central Horticultural Experiment Station farm at Vejalpur, Panchamahals, Gujarat. The results revealed that the crop receiving poultry manure gave significantly higher yields (27.54 t/ha) in okra compared to the other three manures like vermi-compost (23.37 t/ha), farm yard manure (20.43 t/ha), sheep manure (15.77 t/ha) and control (8.82 t/ha). However, between two sources of bio fertilizers *Azotobacter chroococcum* proved better and recorded higher yield (23.56 t/ha) compared to *Azospirillum brasilense* (19.99 t/ha) with an increase of 17.86 percent over the other. Maximum nitrogen (226.2 kg/ha), phosphorous (18.9 kg/ha), potassium (62.4 kg/ha), calcium (157.8 kg/ha), magnesium (129.8 kg/ha) and sulphur (32.1 kg/ha) uptake was observed in the treatment received poultry manure in combination with *Azotobacter chroococcum*.

Published in : HortFlora Research Spectrum, 6 (2) : 114-118 (June 2017)

11. Performance of Tuberose (*Polianthes tuberosa* L.) Cultivars grown for Spikes under Western Uttar Pradesh conditions

Virendra Pal^{1*}, K.P. Singh², Ashish Tyagi¹, Anant Kumar³ and Omvir Singh¹

¹Krishi Vigyan Kendra, Hastinapur,

²Division of Floriculture and Landscaping, Indian Institute of Agricultural Research, Pusa, New Delhi-110012

³Krishi Vigyan Kendra, Muradnagar, Ghaziabad S.V.P.U.A & T, Meerut-250 110 (U.P.)

*Corresponding Author's E-mail: dypgangwar77@gmail.com

ABSTRACT : A field experiment was conducted to assess the performance of tuberose cultivars used for spikes production under Western Uttar Pradesh conditions at Research Farm, under Crop Cafeteria Unit, Krishi Vigyan Kendra, Hastinapur, SVP University of Agriculture & Technology, Meerut (UP) India during the year 2014-15 and 2015-16. Eight potential cultivars namely Mexican Single, Arka Nirantara, Shringar, Rajat Rekha, Pearl Double, Suvasini, Vaibhav and Swarn Rekha were performed in a Randomized Complete Block Design (RCBD) with three replications. Tuberose cultivars varied significantly for growth and flowering parameters. Among the various cultivars tested, the longest spike was recorded in the cv. Shringar (74.09 cm) which was at par with cvs. Mexican Single (73.53 cm), Arka Nirantara (71.24 cm) and Rajat Rekha (71.03 cm), whereas the cultivar Swarn Rekha exhibited shortest spike length (63.86 cm). Number of florets/spike was

found to be maximum in Vaibhav (32.32) being at par with Suvasini (31.32), Swarn Rekha (28.75) and Pearl Double (28.58). The minimum number of florets/spike was observed in cv. Rajat Rekha (17.55). The maximum production of spikes/ha was recorded in Suvasini (184110 Nos.) followed by Vaibhav (183731 Nos.), Shringar (183219 Nos.) and Arka Nirantara (182271 Nos.). The minimum number of spikes/ha was recorded in cv. Swarn Rekha (130640 Nos.).

Published in : HortFlora Research Spectrum, 6 (2) : 119-121 (June 2017)

12. Fruit Production and Biochemical aspects of Seeds of *Euryale ferox* Salisb. under *ex-situ* Conditions

Amit Kumar¹, I.S. Singh^{2*}, Ramesh Kumar¹, V. N. Jha³, A. K. Thakur² and Anil Kumar⁴

¹Department of Biotechnology, LNMU, Darbhanga, Bihar

²ICAR Research Complex for Eastern Region, Research Centre for Makhana, Darbhanga-846 005

³Department of Botany, MRM College, LNMU, Darbhanga, Bihar

⁴Shri Bhola Paswan Shastri Agricultural College, Purnia, BAU, Sabour, Bihar

*Corresponding Author's E-mail: induciah@rediffmail.com

ABSTRACT: Field studies on *Euryale ferox* Salisb. using fresh mature seeds were carried out in experimental plots of Research Centre for Makhana, Darbhanga. Production of fresh seeds has been estimated to be 3.0 t/ha. Biochemical analysis of fresh seeds revealed 55.16% available carbohydrate, 25.73% protein, 12.80% moisture, 4.68% fibre, 1.08% ash and 0.52% fat. The seeds were found to contain amylose 20.22%, P 65.41 mg/100 g, K 36 mg/100 g, Na 47.32 mg/100g, Fe 0.96 mg/100g, Mn 0.92 mg/100g, Cu 0.26 mg 100 g and Zn 0.83 mg/100 g.

Published in : HortFlora Research Spectrum, 6 (2) : 122-124 (June 2017)

13. Effect of Stage of Maturity and Method of Drying on Bio-chemical and Organoleptic Attributes of Ber (*Ziziphus mauritiana* Lamk.) Fruits

V.K. Asati^{1*}, K. P Singh¹ and A. K. Goswami²

¹Department of Horticulture, A. S. (P.G.) College, Lakhaoti, Distt. Bulandshahr, C.C.S. University Meerut (U.P.)

²Div. of Fruits & Horticultural Technology, IARI, New Delhi

*Corresponding author's E-mail: drvinodasati@gmail.com

ABSTRACT : The study was carried out to evaluate the effect of different stages of ripening on the quality of *Z. mauritiana* fruits during drying. The fruits were graded into Immature (Greenish), Mature (Golden Yellow) and Ripe (Redish Brown) categories and these formed the treatments. The dehydration method was more effective than the open sun drying method for retention of fruits colour, reducing fruit spoilage percentage and moisture content during drying. Organoleptic rating was slightly higher in dehydrated fruits than sun drying fruits. In sun drying method per cent acidity was found higher whereas the ascorbic acid content was completely loosed.

Published in : HortFlora Research Spectrum, 6 (2) : 125-127 (June 2017)

14. Response of Bulb size on Vegetative Growth, Flowering and Bulb Production in Tuberose (*Polianthes Tuberosa* L.) cv. Phule Rajani

Krishan P. Singh^{1*}, Tarak N. Saha and P. Hollajer

ICAR-Directorate of Floricultural Research , College of Agriculture Campus, Shivajinagar, Pune – 411005 (Maharashtra)

¹Present address: Division of Floriculture and Landscaping, ICAR-Indian Agricultural Research Institute, New Delhi – 110012

*Corresponding Author's E-mail : amit14_0485@yahoo.com

ABSTRACT : Apart from other production technologies, the performance of tuberose is also influenced by bulb and bulblet to be planted as propagation material. In the present study, beside mother bulb, four bulb grades (diameter in cm) namely, above 3.5 cm, 3.4-2.5, 2.4-1.5, and below 1.5 were selected as treatments for planting. The results revealed that planting of bulb grade above 3.5 cm diameter produced significantly maximum number of leaves/clump (39.15), width of leaf (1.88 cm), number of tillers/clump (14.10), spike length (65.20 cm), rachis length (30.43 cm), fresh weight of cut spike (75.20 cm), number of florets per spike (35.50), number of bulbs per clump (25.90), average diameter of bulb (3.15 cm) and total weight of bulbs/clump (392.40 g). Planting of mother bulbs produced significantly the maximum number of

bulblets per clump (52.35) and their corresponding weight (104.90g). On the other hand, planting of mother bulbs produced significantly the minimum width of leaf (1.46 cm), number of tillers per clump (12.00), rachis length (15.40 cm), fresh weight of cut spike (52.83 g) and weight of bulbs/clump (249.25g). Planting of smallest sized bulbs (below 1.5 cm diameter) produced significantly the minimum number of leaves (17.25) and tillers (12.00) per clump, number of florets/spike (21.18) and number of bulblets (19.05) and their weight (41.65g) per clump.

Published in : HortFlora Research Spectrum, 6 (2) : 128-130 (June 2017)

15. Effect of NPK on Plant Growth and Flowering of *Amaryllis belladonna*

Rohit Kumar Sharma¹, S. Saravanan¹ and Sunita Kumari^{2*}

¹Department of Horticulture, Allahabad School of Agriculture, Sam Higginbottom Institute of Agriculture, Technology and Sciences, Deemed to-be University, Allahabad, INDIA

²Department of Horticulture, G. B. Pant University of Agriculture & Technology, Pantnagar-263 145, Uttarakhand, India

*Corresponding Author's E-mail: sunitakumari.sean@gmail.com

ABSTRACT : The study on effect of NPK on plant growth and flowering of *Amaryllis belladonna* was carried out at Department of Horticulture, Allahabad School of Agriculture, SHIATS, Allahabad during 2013-2014. The study showed that at 180 DAP, application of $N_{125} : P_{50} : K_{125}$ Kg/ha resulted in significantly maximum plant height (43.96 cm) and No. of leaves/plant (8.66), too late flower scape emergence (174.66 days), tallest flower scape (51.76 cm), maximum diameter of flower bud (2.42 cm) and flowers (5.66) and longest durability of flower (6.55 days). The earliest flower scape emergence (167 days) was recorded in $T_1(N_{60} P_{30} K_{30}$ kg/ha) while total number of flower scapes/plant (3.33) and number of flowers/scape (4.10) were found maximum in $T_5 (N_{75} P_{45} K_{45}$ Kg/ha) and $T_5 (N_{175} P_{75} K_{175}$ Kg/ha), respectively.

Published in : HortFlora Research Spectrum, 6 (2) : 131-133 (June 2017)

16. Effect of Mulching on the Growth and Yield of Onion (*Allium cepa* L.)

S.K. Singh¹, Sanjay Kumar² and Subash Chand^{3*}

¹Deptt.of Horticulture, ²Deptt.of Agri. Chemistry and ³Deptt.of Agronomy, B. R. D. (P.G.) College, Deoria

*Corresponding Author's E-mail: subashc7@gmail.com

ABSTRACT : An experiment was conducted at Campus Research Farm, B.R.D. (P.G.) College, Deoria, (U.P.) during the rabi season 2014-15 and 2015-16 to assess the influence of different mulching on the growth and yield of onion. The experiment was laid out in randomized block design with three replications. Treatments of the experiment were seven types of mulching such as control (M_0), Rice straw (M_1), Water hyacinth (M_2), Baggage straw (M_3), Wheat straw (M_4), Grass straw (M_5) and Pipal leaf straw (M_6). The results revealed that growth parameters viz. plant height, no. of leaves, bulb length, bulb diameter, bulb weight and bulb yield were increased significantly with adopting mulching. The maximum plant height, no. of leaves, bulb length, bulb diameter, bulb weight and bulb yield were recorded under Pipal leaf straw. Interestingly, the plant height, no. of leaves, bulb length, bulb diameter, bulb weight and bulb yield did not show significantly differences between M_6 and M_3 . So, mulching with Pipal leaf straw and water hyacinth numerically gave the highest yield 38.00 t/ha, 38.40 t/ha and 37.50 t/ha., 37.60 t/ha during both year of experimentation, respectively.

Published in : HortFlora Research Spectrum, 6 (2) : 134-136 (June 2017)

ICV : 27.39

HORTFLORA RESEARCH SPECTRUM

GIF : 0.471

IBIF : 2.8

NAAS Rating : 3.78

NIIF : 2.14

www.hortflorajournal.com

ISSN : 2250-2823

Published under the Auspices of :

Biosciences and Agriculture Advancement Society (BAAS)

“Shivalay” 98-A Somdutt Vihar, Jagrati Vihar, Garh Road, Meerut-250004

E-mail : hortfloraspectrum.india@gmail.com; submit.hortflorajournal2013@gmail.com



HortFlora Research Spectrum

Quarterly

Published under the Auspices of

BIOSCIENCES & AGRICULTURE ADVANCEMENT SOCIETY (BAAS), Meerut (Regd.)

ISSN : 2250-2823

'Shivalay' 98-A, Somdutt Vihar, Garh Road, Meerut-250 004 (U.P.) India

E-mail : hortfloraspectrum.india@gmail.com, editorhortflora.vku@gmail.com; Mob. : +91 - 9412833903

NAAS Rating : 3.78

Website: www.hortflorajournal.com



Regd.

APPLICATION FORM FOR MEMBERSHIP / SUBSCRIPTION

1. Name (in block letters) :
2. Date of birth :
3. Address for Correspondence :
(in block letters)
- State : PIN :

Photograph

FOR OFFICE USE ONLY

Type of Membership

LM AM IM

Fee Rs.

Receipt No. & Date :

Membership No. : HRS/

Signature of officials

Phone : Fax : E- mail :

4. Occupation: Educationist / Researcher ☐ Instt./ Industry / Business ☐ Student ☐ Others ☐

5. Designation and Official Address :

6. Higher Academic Qualification : Specialization :

7. Professional Experience, if any :

8. Any additional Information :

Type of Membership Desired (tick whichever applicable)

Life membership

(₹ 4000/-)

(US \$ 350)

☐

Annual membership

(₹ 1200/-)

(US \$ 170)

☐

Institutional Membership*

(₹ 2000/-)

(US \$ 250)

☐

Declaration

I wish to become Life / Annual / Institutional Member of the *HortFlora Research Spectrum*. I am enclosing herewith a crossed DD (No..... dated for ₹ issued by in favour of *HortFlora Research Spectrum* payable at Meerut) towards membership/subscription fee of the Journal. If enrolled, I agree to abide by its rules and regulations.

Date :

Place :

Signature :

Journal Subscription Rates (Print Version)

		India	Foreign**
Individual Life Membership	–	₹ 4000/-	US \$ 350
Individual Annual Membership	–	₹ 1200/-	US \$ 170
Library / Corporate Subscription*	–	₹ 2000/-	US \$ 250

*Subscription for one year (One Volume) only. **Only full text PDF.

Duly filled application form along with membership/subscription fee should be mailed to **Managing/Chief Editor, HortFlora Research Spectrum**, 98A, Somdutt Vihar, Garh Road, Meerut - 250 004 (U.P.) India

Membership/subscription fee may also be remitted by Cash at Editorial Office or directly to Journal's Bank Account through e-banking.

Note: Photostat copy of the Application Form may also be used. Each member must submit duly filled application form separately.



ISSN: 2250-2823

HortFlora Research Spectrum

'Shivalay' 98-A, Somdutt Vihar, Garh Road, Meerut-250 004 (U.P.) India
 E-mail : hortfloraspectrum.india@gmail.com, editorhortflora.vku@gmail.com
 Website: www.hortflorajournal.com, Mob. : +91 - 9412833903

GUIDELINES TO THE CONTRIBUTORS & FORMAT FOR ARTICLES

The *HortFlora Research Spectrum*, a Peer Reviewed International Journal, is published Quarterly every year. It publishes original **Review/Strategy Papers, Research Papers and Research Notes** on all facets of Horticulture and allied branches of Science & Technology. The publication is generally open to all Scientists/Researchers/Students of concerned subjects. All the author(s) of the paper must be **Life/Annual** member of the Journal. Duly filled application form for membership/ subscription of the Journal along with prescribed fee should be submitted at the time of submission of manuscript. Each **Life/Annual** member will be given a unique membership number for future reference. Author(s) who are already member/ subscriber of the Journal are requested to quote their Membership No. in covering letter of the manuscript. **Remittance of ₹ 800/- (US\$ 75) per article towards processing & printing charge is mandatory at the time of submission of manuscript.** Membership/subscription fee may be remitted in Cash or through Crossed DD (non-refundable) in favour of *HortFlora Research Spectrum* payable at Meerut. Manuscript typed in MS Word as per the format of the Journal must be submitted via e-mail/online. Hard copy / CD of M/scrip will not be accepted. Authors are also requested to send a Certificate of Originality of paper and No Objection duly signed by all the authors. On receipt of an article at the Editorial Office, an acknowledgement giving the M/scrip number will be sent to the corresponding author which should be quoted while making any future query about its status. All the correspondence regarding membership/ subscription and manuscript submission should be in favour of **Managing/Chief Editor, HortFlora Research Spectrum**, 'Shivalay', 98A, Somdutt Vihar, Garh Road, Meerut-250 004 (U.P.) India.

Format for Manuscript :- Manuscript must be typed (double line space in MS Word, Times New Roman, 12 Font size) on one side of a A4 size paper. References should be properly incorporated in the text along with their serial no. in bracket in place of year. Photos should be in JPG format.
Title of the Paper:- All capitals and bold in 16 pt font (not more than 30 characters)

Author (s):- First letter of name should be Capital & other small letters and bold in 11 pt Times New Roman. If the authors are from different institute(s), they should be properly marked as ^{1, 2, 3}

Full address of institute (Where work actually carried out). E-mail of Corresponding Author

Abstract :- It should be brief, not more than 200 characters in 11pt Font size and 12 lines.

Key words:- Not more than five.

Introduction:- Without heading, 12-15 lines, short, precise, fulfilling objectives of the study.

Materials and Methods :- Heading in capitals, Full details of materials & methods used for experimentation, collection

& analysis of data.

Results and Discussion:- Heading in capitals, Focusing on the fulfilment of stated objectives of the experiment, statistically analysed data presented in the form of tables / figures / photographs. Duplication of data in table and figure should be avoided. Results in form of trends, rather than numerical value should be discussed in the light of authentic available literature. References should properly be incorporated in the text along with serial no. in place of year, e.g. Jayawardena (1), Johnson (2), Kapil and Arora (3), Rashid *et al.* (4) etc. Generic and specific as well as vernacular names should be italicized.

Tables & Figures :- Tables, figures, captions and illustrations should be given in separate sheet properly numbered in Arabic numerals in order of their reference.

Acknowledgement :- If applicable.

References:- In full length papers and in research notes, the number of references should not exceed 15 and 8, respectively. In review/strategy papers it may varies up to 30-40. At the end of the text, references should be arranged alphabetically with proper serial No., Surname first, Year in bracket, Full title of work, Journal name in standard abbreviation and *italic*, Vol No. Bold, Issue No. in bracket, page No. e.g.

1. Jayawardena, S.P. (2013). Effective inoculation method and optimum concentration of *Oryctes* virus for biological control of coconut beetle (*Oryctes rhinoceros*) adults. *HortFlora Res. Spectrum*, 2 (4) : 319-323.
2. Johnson, D.A. (1940). *Plant Microtechnique*, McGraw- Hill Publishing Co. Ltd., New York. PP-29
3. Kapil, R.N. and Arora, S. (1990). Some fascinating features of orchid pollen. *J. Orchid Soc.*, 4 (1): 9-28.
4. Rashid, S., Ashraf, M., Bibi, S. and Anjum, R. (2000). Antibacterial and antifungal activities of *Launaea nudicaulis* Roxb. and *Launaea resedifolia* L. *Pakistan J. Biol. Sci.*, 3 (4) : 630-632.

A full length paper should not exceed 10 pages and a review/strategy article should not exceed 15 pages including tables & figures. In case of review/strategy papers and research notes, the main text is not to have sub headings of Materials & Methods and Results & Discussion. The corresponding author should mention his/her present address with telephone/mobile number and E-mail ID for effective communication.

Acceptance of a manuscript for publication in *HortFlora Research Spectrum* shall automatically mean transfer of copyright to the Journal. The Editorial Board has no responsibility for the statements, opinion or facts expressed in the article published in this Journal, which rests entirely with the Author (s) there of. Editorial Board has also right to format the article as per Journal's format accordingly. PDF file of the published article will be mailed to corresponding author's E-mail for earliest convenience.

Printed & Published by : Dr. Vandana Umrao and **Edited by :** Dr. Vijai Kumar Umrao, Secretary, BAAS
 'Shivalay' 98-A, Somdutt Vihar, Garh Road, Meerut-250 004 (U.P.) INDIA. Mob.: +91-9412833903
 E-mail: hortfloraspectrum.india@gmail.com, editorhortflora.vku@gmail.com
 Website: www.hortflorajournal.com

Printed at : New Rishabh Offset Printers, Delhi Road, Meerut.

ISSN 2250-2823

